
First records of leucism in two species of duiker (Cephalophinae) in Gabon

Stephan Ntie, Graden Froese, Eyeghe Placide Obiang, Joseph Okouyi and Johnny Mfane Otsaghe



Ntie S, Froese G, Obiang EP, Okouyi J, Otsaghe JM 2023. First records of leucism in two species of duiker (Cephalophinae) in Gabon. Biodiversity Observations 13: 233–238.

13 September 2023 (corrected 2023-10-12)

DOI: 10.15641/bo.1390

Mammals

First records of leucism in two species of duiker (Cephalophinae) in Gabon

Stephan Ntie^{1,2*}, Graden Froese³, Eyeghe Placide Obiang⁴, Joseph Okouyi⁵, Johnny Mfane Otsaghe⁶

¹Laboratoire de Biologie Moléculaire et Cellulaire (LABMC), Département de Biologie, Université des Sciences et Techniques de Masuku (USTM), BP 941, Franceville, Gabon

²Cellule Scientifique, Agence Nationale des Parcs Nationaux (ANPN), BP 20379, Libreville, Gabon

³Nsombou Abalghe-Dzal Association (NADA), Makokou, Gabon ⁴Wildlife Conservation Society, Global Conservation Program, New York, USA

⁵Ministère des Eaux et Forêts, Libreville, Gabon ⁶Association Ebyeng-Edzuameniène (A2E), Ebyeng-Edzuameniène, Ogooué-Ivindo Province, Gabon *stephanntie@yahoo.fr

Abstract

The present study aimed to describe partial leucism, which is a pigmentation disorder, in two duiker species: *Philantomba monticola* and *Sylvicapra grimmia*. Leucism is a condition characterized by partial loss of pigmentation, resulting in white or pale patches on the animal's body. This is the first report of leucism in the subfamily Cephalophinae, which includes duikers, and similar occurrences may exist

in other duiker species across their sub-Saharan distribution range. This finding highlights the potential for further research on pigmentation disorders in duikers and their prevalence in different species within the subfamily.

Observation

Leucism (also known as partial albinism) is a pigmentation disorder characterized by a reduction or lack of pigmentation in different parts of the body. It appears in vertebrates as white spots on the skin and/or pure white fur or feather, and normal coloured eyes and claws (Buckley 1982; Miller 2005). Previous studies have shown that it is caused by the expression of mutant alleles (Edelaar et al. 2011; Fleck et al. 2016; Liu et al. 2019). Leucism seems frequent in bats (Marin-Vasquez et al. 2010; Sánchez-Hernández et al. 2012), birds (Sage 1962, 1963; Bensch et al. 2000; van Grouw 2006; Guay et al. 2012; van Grouw 2012) and in other mammals (ie. Carnivores, rodents (Acevedo & Aguayo 2008; Steen & Sonerud 2012; Abreu et al. 2013; Brito & Valdivieso-Bermeo 2016; Talamoni et al. 2017). The goal of this paper was to report leucism in two duiker species: the blue duiker *Philantomba monticola* and the common duiker *Sylvicapra grimmia* (Figure 1).

The first case of leucism in blue duiker was reported in July 2018 in northern Gabon, in the village called "Konoville les deux Eglises" (1°50'28"N 11°38'39"E), in the Woleu-Ntem Province. This specimen of blue duiker was caught on a cable snare trap (Figure 2a), skinned (Figure 2b and c), and the meat sold to locals. The entire body of the animal was white, except for a few areas around the ears, the dorsal part of the neck, and the upper part of the back (in continuity with the neck). Although this part of the country is more populated and has more plantations, primary forest is still present (Wheatley 2014).

The second case of leucism in blue duiker was reported in August 2021 in northeastern Gabon, near the town of Makokou (0°34′00″ N, 12° 52′ 00″E), in the Ivindo Province. There, the blue duiker specimen was white only on the tips of both its hind legs, the rest of body




Figure 1: Location of the three reported cases of duiker leucism in Gabon: (a and b) *Philantomba monticola* and (c) *Sylvicapra grimmia*

being normally coloured (Figure 3). The blue duiker is commonly hunted and sold as bushmeat in the region (Juste et al. 1995; Fa et al. 2000; Barnes 2002; Fa et al. 2002; Nielsen 2006; Kümpel et al. 2010; Ávila et al. 2019). In the wild, this species occupies different habitat types including dense forest, woodland mosaics, and forest fragments (Dubost 1980; Gautierhion et al. 1980; Heymans & Lejoly 1981; Kingdon 1997). This part of the country is covered with dense tropical forest and the Ivindo National Park, a new UNESCO world heritage site, is nearby (UNESCO World Heritage Convention 2021).

For the present study, a third leucistic duiker was reported in August 2021 in southeastern Gabon, within the Leconi Park (1°25' 24.5"S, 14°16' 50.9"E). The species is the common duiker *S. grimmia*. A young duiker was observed close to the leucistic duiker, and was mostly likely its cub. The young had normal body coloration, while the adult looked entirely white from afar (Figure 4). The common duiker is the only one that is exclusively found in savanna habitat

(Kingdon 1997). In Gabon, the common duiker is solely found in the Plateau Batéké region (including in Gabon's Plateau Batéké National Park) where there are extensive savannas which span into the neighbouring country, the Republic of the Congo.

The present study is the first to report cases of leucism in duikers. These observations call into question the species designation of the white-legged duiker *C. Ogilbyi* which is genetically close to the Peter's duiker *C. callypigus* (Ntie et al. 2010; Johnston et al. 2011; Johnston & Anthony 2012; Ghassemi-Khademi & Hamidi 2019).

References

Abreu M, Machado R, Barbieri F, Freitas N, Oliveira L 2013.

Anomalous colour in Neotropical mammals: a review with new records for *Didelphis* sp. (Didelphidae, Didelphimorphia) and *Arctocephalus australis* (Otariidae, Carnivora). Brazilian Journal of Biology 73: 185–194.

Acevedo J, Aguayo M 2008. Leucistic South American sea lion in Chile, with a review of anomalously color in otariids. Revista de Biologia Marina y Oceanografia 43: 413–417.

Ávila E, Tagg N, Willie J, Mbohli D, Farfán M.Á, Vargas JM, Bonat WH, Dupain J, Epanda MA, Luyten I 2019. Interpreting long-term trends in bushmeat harvest in southeast Cameroon. Acta Oecologica 94: 57–65.

Barnes RF 2002. The bushmeat boom and bust in West and Central Africa. Oryx 36: 236–242.

Bensch S, Hansson B, Hasselquist D, Nielsen B 2000. Partial albinism in a semi-isolated population of Great Reed Warblers. Hereditas 133: 167–170.



Figure 2: The blue duiker *Philantomba monticola* specimen reported by the village Konoville les deux Eglises, in the Woleu-Ntem Province, northern Gabon: (a) the entire specimen, and (b and c) and the left-over skin after the meat was sold.



b)

Figure 3: The blue duiker *Philantomba monticola* specimen reported near the city of Makokou, Ogooué Ivindo Province, northeastern Gabon.

Figure 4: The common duiker *Sylvicapra grimmia* (a and b) specimen reported in Lekoni Park, southeastern Gabon.

- **Brito J, Valdivieso-Bermeo K 2016.** First records of leucism in eight species of small mammals (Mammalia: Rodentia). Therya 7: 483 –489.
- **Buckley PA 1982**. Avian genetics. In: Diseases of Cage and Aviary Birds, 2nd Ed. Petrak ML (ed.), pp 21–110. Lea and Febiger, Philadelphia, USA.
- **Dubost G 1980**. Ecology and social-behavior of the Blue Duiker small African forest ruminant. Zeitschrift fur Tierpsychologie/ Journal of Comparative Ethology 54: 205–266.
- Edelaar P, Donázar JA, Soriano M, Santillán MÁ, González-Zevallos D, García Borboroglu P, Lisnizer N, Gatto AJ, Agüero ML, Passera CA 2011. Apparent selective advantage of leucism in a coastal population of Southern Caracaras (Falconidae). Evolutionary Ecology Research 13: 187–196.
- **Fa JE, Juste J, Burn RW, Broad G 2002**. Bushmeat consumption and preferences of two ethnic groups in Bioko Island, West Africa. Human Ecology 30: 397–416.
- **Fa JE, Yuste JEG, Castelo R 2000**. Bushmeat markets on Bioko Island as a measure of hunting pressure. Conservation Biology 14: 1602–1613.
- **Fleck K, Erhardt G, Luehken G 2016**. From single nucleotide substitutions up to chromosomal deletions: genetic pause of leucismassociated disorders in animals. Berliner und Munchener Tierarztliche Wochenschrift 129: 269–281.
- **Gautierhion A, Emmons LH, Dubost G 1980**. A comparison of the diets of three major groups of primary consumers of Gabon (primates, squirrels and ruminants). Oecologia 45: 182–189.
- **Ghassemi-Khademi T, Hamidi K 2019**. Re-evaluation of molecular phylogeny of the Subfamily Cephalophinae (Bovidae: Artiodacty-

- la); with notes on diversification of body size. Jordan Journal of Biological Sciences 12: 637–647.
- **Guay P-J, Potvin DA, Robinson RW 2012**. Aberrations in plumage coloration in birds. Australian Field Ornithology 29: 23–30.
- Heymans J, Lejoly J. 1981. Sur la répartition des Cephalophinae en forêt ombrophile (Région du Haut-Zaïre, Rép. du Zaïre) (Bovidae). Revue de Zoologie Africaine 95: 1–10.
- **Johnston AR, Anthony NM 2012**. A multi-locus species phylogeny of African forest duikers in the subfamily Cephalophinae: evidence for a recent radiation in the Pleistocene. BMC Evolutionary Biology 12: 120.
- Johnston AR, Morikawa MK, Ntie S, Anthony NM 2011. Evaluating DNA barcoding criteria using African duiker antelope (Cephalophinae) as a test case. Conservation Genetics 12: 1173.
- **Juste J, Fa JE, Del Val JP, Castroviejo J. 1995**. Market dynamics of bushmeat species in Equatorial Guinea. Journal of Applied Ecology: 454–467.
- **Kingdon J. 1997**. The Kingdon field guide to African mammals. Academic Press, London and New York.
- Kümpel NF, Milner-Gulland EJ, Cowlishaw G, Rowcliffe JM 2010.

 Assessing sustainability at multiple scales in a rotational bushmeat hunting system. Conservation Biology 24: 861–871.
- Liu Z, Wen H, Hailer F, Dong F, Yang Z, Liu T, Han L, Shi F, Hu Y, Zhou J 2019. Pseudogenization of Mc1r gene associated with transcriptional changes related to melanogenesis explains leucistic phenotypes in *Oreonectes* cavefish (Cypriniformes, Nemacheilidae). Journal of Zoological Systematics and Evolutionary Research 57: 900–909.

- Marin-Vasquez A, Ortega-Rincón M, Ramírez-Chaves HE 2010.
 Records of leucism in three species of Colombian bats: Carollia brevicauda, Artibeus jamaicensis and Lophostoma silvicolum (Phyllostomidae). Chiroptera Neotropical 16: 706–709.
- Miller J 2005. All about albinism. Missouri Conservationist 66: 4–7.
- **Nielsen MR 2006**. Importance, cause and effect of bushmeat hunting in the Udzungwa Mountains, Tanzania: Implications for community-based wildlife management. Biological Conservation 128: 509–516.
- Ntie S, Johnston AR, Mickala P, Bowkett AE, van Vuuren BJ, Colyn M, Telfer P, Maisels F, Hymas O, Rouyer RL, Wallace RA, LeBlanc K, van Vliet N, Sonet G, Verheyen E, Pires D, Wickings EJ, Lahm SA, Anthony NM 2010. A molecular diagnostic for identifying central African forest artiodactyls from faecal pellets. Animal Conservation 13: 80–93.
- **Sage BL 1962**. Albinism and melanism in birds. British Birds 55: 201–225.
- **Sage BL 1963**. The incidence of albinism and melanism in British birds. British Birds 56: 409–416.
- Sánchez-Hernández C, Rojas-Martínez A, López-Vidal JC, Elizalde-Arellano C, Romero-Almaraz M.d.L, Aguillar-López M, Taboada-Salgado A 2012. Leucism in five species of bats from Mexico. Chiroptera Neotropical 18: 1123–1127.
- **Steen R, Sonerud GA 2012**. A bank vole (*Myodes glareolus*) with complete leucism captured by a Eurasian kestrel (*Falco tinnunculus*) in Norway. Annales Zoologici Fennici 49: 306–308.
- Talamoni S, Viana PIM, Costa CG, Palú L, Oliveira RB, Pessôa LM 2017. Occurrence of leucism in *Eira barbara* (Carnivora, Mustelidae) in Brazil. Biota Neotropica. 17(3): e20170328. Avail-

able online at https://www.biotaneotropica.org.br/BN/article/view/1456

- **UNESCO World Heritage Convention 2021**. Ivindo National Park. Available online at https://whc.unesco.org/en/list/1653/
- van Grouw H 2006. Not every white bird is an albino: sense and nonsense about colour aberrations in birds. Dutch Birding 28: 79– 89.

tion/291264948 What Colour Is That Sparrow A Case Study Colour Aberrations In The House Sparrow Passer Domes ticus

Wheatley N 2014. Where to Watch Birds in Africa (Gabon pp. 139–152). Princeton University Press, Princton, USA.

Paper edited by Megan Loftie-Eaton Biodiversity and Development Institute



Biodiversity Observations is powered by Open Journal Systems (OJS) and is hosted by the University of Cape Town Libraries. OJS is an open source software application for managing and publishing scholarly journals. Developed and released by the Public Knowledge Project in 2001, it is the most widely used open source journal publishing platform in existence, with over 30,000 journals using it worldwide.